

PRODUCT INFORMATION

Target CLIC3
Synonyms N/A

Description Human CLIC3 full length protein-synthetic nanodisc

Delivery 6~8weeks
Uniprot ID 095833
Expression Host HEK293

Protein Families Ion Channels: Other

Protein Pathways N/A

Background

Molecular Weight The human full length CLIC3 protein has a MW of

26.6kDa

Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5%

- 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with

pH lower than 6.5 in subsequent experiments.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store

Storage & Shipping Intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient

temperature.

Chloride channels are a diverse group of proteins that regulate fundamental cellular processes including stabilization of cell membrane potential, transepithelial transport, maintenance of

transepithelial transport, maintenance of intracellular pH, and regulation of cell volume. Chloride intracellular channel 3 is a member of the p64 family and is predominantly localized in

the p64 family and is predominantly localized in the nucleus and stimulates chloride ion channel activity. In addition, this protein may participate in cellular growth control, based on its association with ERK7, a member of the MAP kinase family.

> Email: info@dimabio.com Website: www.dimabio.com

[provided by RefSeq, Jul 2008]

Usage Research use only

