

**PRODUCT INFORMATION**

<b>Target</b>	CMTM6
<b>Synonyms</b>	CKLFSF6; PRO2219
<b>Description</b>	Human CMTM6 full length protein-synthetic nanodisc
<b>Delivery</b>	3-4 weeks
<b>Uniprot ID</b>	Q9NX76
<b>Expression Host</b>	HEK293
<b>Protein Families</b>	Transmembrane
<b>Protein Pathways</b>	N/A
<b>Molecular Weight</b>	The human full length CMTM6 protein has a MW of 20.4 kDa
<b>Formulation &amp; Reconstitution</b>	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with pH lower than 6.5 in subsequent experiments.
<b>Storage &amp; Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	This gene belongs to the chemokine-like factor gene superfamily, a novel family that is similar to the chemokine and transmembrane 4 superfamilies. This gene is one of several chemokine-like factor genes located in a cluster on chromosome 3. This gene is widely expressed in many tissues, but the exact function of the encoded protein is unknown. [provided by RefSeq, Jul 2008]
<b>Usage</b>	Research use only



### ELISA assay to evaluate CMTM6-Nanodisc 0.2 $\mu$ g Human CMTM6-Nanodisc per well




Figure1. Elisa plates were pre-coated with Flag Tag CMTM6-Nanodisc (0.2 $\mu$ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with CMTM6-Nanodisc is 4.393ng/ml.



Figure2. Human CMTM6-Nanodisc, Flag Tag on SDS-PAGE

