

PRODUCT INFORMATION

Target	CXCL6
Synonyms	C-X-C Motif Chemokine 6; Chemokine Alpha 3; CKA-3; Granulocyte Chemotactic Protein 2; GCP-2; Small-Inducible Cytokine B6; CXCL6; GCP2; SCYB6
Description	Recombinant Human C-X-C Motif Chemokine 6 is produced by our Mammalian expression system and the target gene encoding Gly38-Asn114 is expressed with a 6His tag at the C-terminus.
Delivery	In Stock
Uniprot ID	P80162
Expression Host	HEK293
Tag	
Molecular Characterization	Not available
Molecular Weight	9.35 KDa
Purity	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
Formulation & Reconstitution	Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, 5% Trehalose, 1mM EDTA, pH 7.4.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	Chemokine (C-X-C-Motif) Ligand 6 (CXCL6) is a small cytokine belonging to the CXC chemokine family. It is a potent neutrophil chemotactic and activating factor and it exhibits extensive similarity to other CXC chemokines such as IL-8 and ENA-78. CXCL6 can promote the release of MMP-9 from granulocytes indicating its potential role as an inflammatory mediator. It functionally uses both of the IL-8/CXCL8 receptors to chemoattract neutrophils but that is structurally most related to epithelial cell-derived neutrophil attractant-78 (ENA-78)/CXCL5. The human CXCL6 gene has been cloned and is physically mapped to the CXC chemokine locus on chromosome 4. Mature human CXCL6 is a 75 amino acid (aa) protein with a predicted molecular weight of approximately 8 kDa. Human CXCL6 shares 60% and 67% aa identity with mouse and bovine CXCL6, respectively.
Usage	Research use only
Conjugate	Unconjugated






Figure 1. Greater than 95% as determined by reducing SDS-PAGE.

