

PRODUCT INFORMATION

| | |
|------------------------------|--|
| Tag | C-Flag Tag |
| Target | CD39 |
| Synonyms | ATPDase;ENTPD1;NTPDase-1;SPG64 |
| Description | Human CD39 full length protein-synthetic nanodisc |
| Delivery | In Stock |
| Uniprot ID | P49961 |
| Expression Host | HEK293 |
| Protein Families | Transmembrane |
| Protein Pathways | Purine metabolism, Pyrimidine metabolism |
| Molecular Weight | The human full length CD39 protein has a MW of 58.0 kDa |
| Formulation & Reconstitution | Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments. |
| Storage & Shipping | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Background | The protein is a plasma membrane protein that hydrolyzes extracellular ATP and ADP to AMP. Inhibition of this protein's activity may confer anticancer benefits. Several transcript variants encoding different isoforms have been found for this gene. |
| Usage | Research use only |
| Conjugate | Unconjugated |






Figure1. Elisa plates were pre-coated with Flag Tag CD39-Nanodisc (0.2µg/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with CD39-Nanodisc is 7.763ng/ml.




Figure2. Human CD39-Nanodisc, Flag Tag on SDS-PAGE

