

## PRODUCT INFORMATION

<b>Target</b>	EREG
<b>Synonyms</b>	Ep;EPR;ER
<b>Description</b>	Recombinant Human EREG with C-terminal human Fc tag
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	O14944
<b>Expression Host</b>	HEK293
<b>Tag</b>	C-Human Fc Tag
<b>Molecular Characterization</b>	EREG(Val63-Leu108) hFc(Glu99-Ala330)
<b>Molecular Weight</b>	The protein has a predicted molecular mass of 31.4 kDa after removal of the signal peptide. The apparent molecular mass of EREG-hFc is approximately 35-55 kDa due to glycosylation.
<b>Purity</b>	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Storage &amp; Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	This gene encodes a secreted peptide hormone and member of the epidermal growth factor (EGF) family of proteins. The encoded protein is a ligand of the epidermal growth factor receptor (EGFR) and the structurally related erb-b2 receptor tyrosine kinase 4 (ERBB4). The encoded protein may be involved in a wide range of biological processes including inflammation, wound healing, oocyte maturation, and cell proliferation. Additionally, the encoded protein may promote the progression of cancers of various human tissues. [provided by RefSeq, Jul 2015]
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated






Figure 1. Human EREG Protein, hFc Tag on SDS-PAGE under reducing condition.

### Human EREG, hFc Tagged protein ELISA

0.2 µg of Human EREG, hFc tagged protein per well




Figure 2. ELISA plate pre-coated by 2 µg/mL (100 µL/well) Human EREG Protein, hFc Tag (PME100617) can bind Anti-EREG antibody (DMC424), IgG1 Chimeric mAb in a linear range of 0.64-16 ng/mL.

