

PRODUCT INFORMATION

Tag C-Flag&Strep Tag

Target F2RL1

Synonyms GPR11; PAR2

DescriptionHuman F2RL1-Strep full length protein-synthetic

nanodisc

Delivery 6~8weeks

Uniprot ID P55085

Expression Host HEK293

Protein Families Druggable Genome, GPCR, Transmembrane

Protein Pathways Neuroactive ligand-receptor interaction

Molecular Weight

The human full length F2RL1-Strep protein has a

MW of 44.1 kDa Lyophilized from nanodisc solubilization buffer (20

mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5%
– 8% trehalose is added as protectants before
lyophilization. Please see Certificate of Analysis

Formulation & Reconstitutionlyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high

concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store

Storage & Shipping Interlued for use within a month, and obtained and store at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient

temperature.

A member of the G-protein coupled receptor 1 family of proteins. The encoded cell surface receptor is activated through proteolytic cleavage of its extracellular amino terminus, resulting in a new amino terminus that acts as a tethered ligand that binds to an extracellular loop domain. Activation of the receptor has been shown to stimulate vascular smooth muscle relaxation,

stimulate vascular smooth muscle relaxation, dilate blood vessels, increase blood flow, and lower blood pressure. This protein is also important in the inflammatory response, as well

Email: info@dimabio.com Website: www.dimabio.com

as innate and adaptive immunity.

Usage Research use only
Conjugate Unconjugated

Background

