

PRODUCT INFORMATION

FCRL5 **Target**

CD307; FCRH5; IRTA2; BXMAS1; CD307e; PRO820 **Synonyms**

Recombinant human FCRL5(745-847)Protein with **Description**

C-terminal mouse Fc tag

Delivery In Stock **Uniprot ID Q96RD9 Expression Host HEK293**

Tag C-mouse Fc tag

Molecular

Background

FCRL5(Val745-Pro847)mFc(Pro99-Lys330) Characterization

The protein has a predicted molecular mass of **Molecular Weight**

37.0 kDa after removal of the signal peptide. The apparent molecular mass of FCRL5(745-847)-mFc is approximately 35-55 kDa due to glycosylation. The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue

Purity

staining.

Lyophilized from sterile PBS, pH 7.4. Normally 5 % 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Formulation & Reconstitution

for specific instructions.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store Storage & Shipping

at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

This gene encodes a member of the immunoglobulin receptor superfamily and the Fc-

receptor like family. This gene and several other Fc receptor-like gene members are clustered on the long arm of chromosome 1. The encoded protein is a single-pass type I membrane protein and contains 8 immunoglobulin-like C2-type

domains. This gene is implicated in B cell development and lymphomagenesis. Alternatively spliced transcript variants encoding different isoforms have been identified. [provided by RefSeq, Sep 2010]

> Email: info@dimabio.com Website: www.dimabio.com

Usage Research use only

Conjugate Unconjugated



Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China)

/+86-400-006-0995(China)









Figure 1. Human FCRL5(745-847) Protein, mFc Tag on SDS-PAGE under reducing condition.

Human FCRL5(745-847), mFc Tagged protein ELISA

0.2 μg of Human FCRL5(745-847), mFc tagged protein per well



Anti-FCRL5(cevostamab biosimilar without anti-CD3 portion) mAb (ng/mL)

Figure 2. ELISA plate pre-coated by 2 μ g/mL (100 μ L/well) Human FCRL5(745-847) Protein, mFc Tag (PME101495) can bind Anti-FCRL5(cevostamab biosimilar without anti-CD3 portion) mAb (BME100089) in a linear range of 0.64–80 ng/mL.

Email: info@dimabio.com Website: www.dimabio.com

