

## PRODUCT INFORMATION

<b>Target</b>	FGF-19
<b>Synonyms</b>	Fibroblast growth factor 19;FGF-19;FGF19
<b>Description</b>	Recombinant Human Fibroblast Growth Factor 19 is produced by our E.coli expression system and the target gene encoding Phe27-Lys216 is expressed with a 6His tag at the N-terminus.
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	O95750
<b>Expression Host</b>	E.coli
<b>Tag</b>	
<b>Molecular Characterization</b>	Not available
<b>Molecular Weight</b>	23.5 KDa
<b>Purity</b>	Greater than 95% as determined by reducing SDS-PAGE.
<b>Formulation &amp; Reconstitution</b>	Lyophilized from a 0.2 µm filtered solution of 20mM Tris-HCl, 150mM NaCl, 1mM EDTA, pH 8.0.
<b>Storage &amp; Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	Fibroblast growth factor 19 (FGF19) is a secreted protein which belongs to the FGFs family. FGF19 is expressed in fetal brain, cartilage, retina, and adult gall bladder. FGFs modulate cellular activity via at least 5 distinct subfamilies of high-affinity FGF receptors (FGFRs): FGFR-1, -2, -3, and -4, all with intrinsic tyrosine kinase activity. FGFRs can be important for regulation of glucose and lipid homeostasis. FGF19 has important roles as a hormone produced in the ileum in response to bile acid absorption. It has been shown to cause resistance to diet-induced obesity and insulin desensitization and to improve insulin, glucose, and lipid profiles in diabetic rodents. FGF19 can be considered as a regulator of energy expenditure.
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated






Figure 1. Greater than 95% as determined by reducing SDS-PAGE.

DIMABIO CONFIDENTIAL

