Cat. No. FLP100130



## **PRODUCT INFORMATION**

C-Flag Tag Tag

**GIPR Target Synonyms** PGOTL2

Human GIPR full length protein-synthetic **Description** 

nanodisc **Delivery** In Stock **Uniprot ID** P48546 **Expression Host HEK293** 

**Protein Families** Druggable Genome, GPCR, Transmembrane

**Protein Pathways** Neuroactive ligand-receptor interaction

The human full length GIPR protein has a MW of **Molecular Weight** 

53.2 kDa Lyophilized from nanodisc solubilization buffer (20

mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Formulation &

Reconstitution for specific instructions. Do not use solvents with a pH below 6.5 or those containing high

concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store

Storage & Shipping at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

témperature.

A G-protein coupled receptor for gastric inhibitory polypeptide (GIP), which was originally identified as an activity in gut extracts that inhibited gastric

acid secretion and gastrin release, but subsequently was demonstrated to stimulate

**Background** insulin release in the presence of elevated

glucose. Mice lacking this gene exhibit higher blood glucose levels with impaired initial insulin response after oral glucose load. Defect in this gene thus may contribute to the pathogenesis of

> Email: info@dimabio.com Website: www.dimabio.com

diabetes.

Usage Research use only

Conjugate Unconjugated

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China)

/+86-400-006-0995(China)





## **ELISA assay to evaluate GIPR-Nanodisc** 0.2µg Human GIPR-Nanodisc per well




Figure 1. Elisa plates were pre-coated with Flag Tag GIPR-Nanodisc (0.2 $\mu$ g/per well). Serial diluted anti-GIPR monoclonal antibody (BME100209) solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-GIPR monoclonal antibody binding with GIPR-Nanodisc is 862.5 $\mu$ g/ml.




Figure 2. Human GIPR-Nanodisc, Flag Tag on SDS-PAGE

Email: info@dimabio.com Website: www.dimabio.com

