## PRODUCT INFORMATION

| | |
|---|---|
| **Target** | GITR |
| **Synonyms** | AITR;GITR;TNFRSF18;CD357 |
| **Description** | Recombinant human GITR protein with C-terminal mouse Fc and 6×His tag |
| **Delivery** | In Stock |
| **Uniprot ID** | Q9Y5U5 |
| **Expression Host** | HEK293 |
| **Tag** | C-Mouse Fc and 6×His Tag |
| **Molecular Characterization** | GITR(Gln26-Pro162) mFc(Pro99-Lys330) 6×His |
| **Molecular Weight** | The protein has a predicted molecular mass of 50-52 kDa after removal of the signal peptide. |
| **Purity** | The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining. |
| **Formulation & Reconstitution** | Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. |
| **Storage & Shipping** | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| **Background** | This gene encodes a member of the TNF-receptor superfamily. The encoded receptor has been shown to have increased expression upon T-cell activation, and it is thought to play a key role in dominant immunological self-tolerance maintained by CD25( )CD4( ) regulatory T cells. Knockout studies in mice also suggest the role of this receptor is in the regulation of CD3-driven T-cell activation and programmed cell death. Three alternatively spliced transcript variants of this gene encoding distinct isoforms have been reported. |
| **Usage** | Research use only |
| **Conjugate** | Unconjugated |



Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China
Telephone: +1 2409940618(USA)
/+86-18062749453(China)
/+86-400-006-0995(China)

Email: info@dimabio.com
Website: www.dimabio.com




Figure 1. Human GITR Protein, mFc-His Tag on SDS-PAGE under reducing condition.




Figure 2. HEK293 cell line transfected with irrelevant protein (B) and human GITR Ligand (A) were surface stained with Human GITR, mFc-His tagged protein (PME100019) 1μg/ml followed by Alexa 488-conjugated anti-mouse IgG secondary antibody.



Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China
Telephone: +1 2409940618(USA)
/+86-18062749453(China)
/+86-400-006-0995(China)

Email: info@dimabio.com
Website: www.dimabio.com