

PRODUCT INFORMATION

| | |
|---|--|
| Tag | C-Flag Tag |
| Target | GPR62 |
| Synonyms | GPCR8, KPG_005 |
| Description | Human GPR62 full length protein-synthetic nanodisc |
| Delivery | 6~8weeks |
| Uniprot ID | Q9BZJ7 |
| Expression Host | HEK293 |
| Protein Families | GPCR,Transmembrane,Druggable Genome, |
| Protein Pathways | GPCRDB Other, |
| Molecular Weight | The human full length GPR62 protein has a MW of 37.6kDa |
| Formulation & Reconstitution | Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments. |
| Storage & Shipping | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Background | Orphan G-protein coupled receptor. Constitutively activates the G(q/11)/inositol phosphate and the G(s)-alpha/cAMP signaling pathways (PubMed:28827538). Has spontaneous activity for beta-arrestin recruitment (PubMed:28827538). Shows a reciprocal modulation of signaling functions with the melatonin receptor MTNR1B most likely through receptor heteromerization (PubMed:28827538).[UniProtKB/Swiss-Prot Function] |
| Usage | Research use only |
| Conjugate | Unconjugated |

