

PRODUCT INFORMATION

Tag	C-Flag Tag
Target	GRIK1
Synonyms	EAA3, EEA3, GLR5, GLUR5, GluK1, gluR-5
Description	Human GRIK1 full length protein-synthetic nanodisc
Delivery	6~8weeks
Uniprot ID	P39086
Expression Host	HEK293
Protein Families	Ion Channels: Glutamate Receptors
Protein Pathways	N/A
Molecular Weight	The human full length GRIK1 protein has a MW of 104kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	Glutamate receptors are the predominant excitatory neurotransmitter receptors in the mammalian brain and are activated in a variety of normal neurophysiologic processes. This gene product belongs to the kainate family of glutamate receptors, which are composed of four subunits and function as ligand-activated ion channels. The subunit encoded by this gene is subject to RNA editing (CAG->CGG; Q->R) within the second transmembrane domain, which is thought to alter the properties of ion flow. Alternative splicing, resulting in transcript variants encoding different isoforms, has been noted for this gene. [provided by RefSeq, Jul 2008]
Usage	Research use only
Conjugate	Unconjugated

