

PRODUCT INFORMATION

Tag C-Flag Tag

Target HCN1

Synonyms BCNG-1, BCNG1, DEE24, EIEE24, GEFSP10, HAC-2

DescriptionHuman HCN1 full length protein-synthetic

Delivery 6~8weeks
Uniprot ID 060741
Expression Host HEK293

Protein Families Ion Channels: Cyclic nucleotide gated

Protein Pathways N/A

Molecular Weight

The human full length HCN1 protein has a MW of

98.8kDa
Lyophilized from nanodisc solubilization buffer (20

mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5%
- 8% trehalose is added as protectants before
lyophilization. Please see Certificate of Analysis
for specific instructions. Do not use solvents with

Reconstitutionfor specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments.

than 5 mM) in subsequent metal lons (greater than 5 mM) in subsequent experiments. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store

Storage & Shipping

Interlided for use within a month, and out and store at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient

témperature.

The membrane protein encoded by this gene is a hyperpolarization-activated cation channel that contributes to the native pacemaker currents in heart and neurons. The encoded protein can homodimerize or heterodimerize with other poreforming subunits to form a potassium channel.

Email: info@dimabio.com Website: www.dimabio.com

This channel may act as a receptor for sour tastes. [provided by RefSeq, Oct 2011]

Usage Research use only

Conjugate Unconjugated

Background



