

## **PRODUCT INFORMATION**

| Тад                             | C-Flag Tag  |
|---------------------------------|---|
| Target                          | HCN3  |
| Synonyms                        | N/A   |
| Description                     | Human HCN3 full length protein-synthetic<br>nanodisc  |
| Delivery                        | 6~8weeks  |
| Uniprot ID                      | Q9P1Z3  |
| <b>Expression Host</b>          | HEK293  |
| <b>Protein Families</b>         | Ion Channels: Cyclic nucleotide gated   |
| Protein Pathways                | N/A   |
| Molecular Weight                | The human full length HCN3 protein has a MW of 86kDa  |
| Formulation &<br>Reconstitution | Lyophilized from nanodisc solubilization buffer (20<br>mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5%<br>– 8% trehalose is added as protectants before<br>lyophilization. Please see Certificate of Analysis<br>for specific instructions. Do not use solvents with<br>a pH below 6.5 or those containing high<br>concentrations of divalent metal ions (greater<br>than 5 mM) in subsequent experiments. |
| Storage & Shipping              | Store at -20°C to -80°C for 12 months in<br>lyophilized form. After reconstitution, if not<br>intended for use within a month, aliquot and store<br>at -80°C (Avoid repeated freezing and thawing).<br>Lyophilized proteins are shipped at ambient<br>temperature.  |
| Background                      | This gene encodes a multi-pass membrane<br>protein that functions as a voltage gated cation<br>channel. The encoded protein is a member of a<br>family of closely related cyclic adenosine<br>monophosphate-binding channel proteins.<br>Alternative splicing results in multiple transcript<br>variants. [provided by RefSeq, Aug 2012]  |
| Usage                           | Research use only   |
| Conjugate                       | Unconjugated  |
|                                 |   |

Email: info@dimabio.com Website: www.dimabio.com

