

## **PRODUCT INFORMATION**

**Target** HLA-A

**Synonyms** Aw-33; Aw-74; HLAA

Recombinant human HLA-A protein with C-Description

terminal human Fc tag

**Delivery** Under development

**Uniprot ID** P04439 **Expression Host HEK293** 

Tag C-Human Fc Tag

Molecular

**Purity** 

**Background** 

HLA-A(Gly25-Ile308) hFc(Glu99-Ala330) Characterization

The protein has a predicted molecular mass of **Molecular Weight** 56.76 kDa after removal of the signal peptide.

The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue

staining.

Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before Formulation & Reconstitution

lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in

lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).

Storage & Shipping

Lyophilized proteins are shipped at ambient

temperature.

HLA-A belongs to the HLA class I heavy chain paralogues. This class I molecule is a heterodimer consisting of a heavy chain and a light chain (beta-2 microglobulin). The heavy chain is anchored in the membrane. Class I molecules play a central role in the immune system by

pray a Central fole in the immune system by presenting peptides derived from the endoplasmic reticulum lumen. They are expressed in nearly all cells. The heavy chain is approximately 45 kDa and its gene contains 8 exons. Exon 1 encodes the leader peptide, exons 2 and 3 encode the alpha1 and alpha2 domains, which both bind the portide.

which both bind the peptide, exon 4 encodes the alpha3 domain, exon 5 encodes the transmembrane region, and exons 6 and 7 encode the cytoplasmic tail. Polymorphisms within exon 2 and exon 3 are responsible for the peptide binding specificity of each class one molecule. Typing for these polymorphisms is continuous for home marrow and kidney. routinely done for bone marrow and kidney

transplantation. Hundreds of HLA-A alleles have been described.

Usage Research use only

Conjugate Unconjugated

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China)

/+86-400-006-0995(China)

Email: info@dimabio.com Website: www.dimabio.com

