

## **PRODUCT INFORMATION**

C-Flag Tag Tag

**Target** HRH3

**Synonyms** GPCR97; HH3R

Human HRH3 full length protein-synthetic **Description** 

nanodisc **Delivery** In Stock **Uniprot ID** Q9Y5N1 **Expression Host HEK293** 

**Protein Families** Druggable Genome, Transmembrane **Protein Pathways** Neuroactive ligand-receptor interaction

The human full length HRH3 protein has a MW of **Molecular Weight** 

48.7 kDa Lyophilized from nanodisc solubilization buffer (20

mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Formulation & Reconstitution

for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater

than 5 mM) in subsequent experiments. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store Storage & Shipping at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

témperature.

Histamine is a ubiquitous messenger molecule released from mast cells, enterochromaffin-like cells, and neurons. Its various actions are mediated by histamine receptors H1, H2, H3 and H4. This gene encodes one of the histamine receptors (H3) which belongs to the family 1 of G protein-coupled receptors. It is an integral

membrane protein and can regulate

neurotransmitter release. This receptor can also increase voltage-dependent calcium current in smooth muscles and innervates the blood vessels

> Email: info@dimabio.com Website: www.dimabio.com

and the heart in cardiovascular system.

Usage Research use only

Conjugate Unconjugated

**Background** 





## ELISA assay to evaluate HRH3-Nanodisc 0.2µg Human HRH3-Nanodisc per well




Figure 1. Elisa plates were pre-coated with Flag Tag HRH3-Nanodisc (0.2 $\mu$ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with HRH3-Nanodisc is 2.708ng/ml.

Email: info@dimabio.com Website: www.dimabio.com



Figure 2. Human HRH3-Nanodisc, Flag Tag on SDS-PAGE

