Human IFNB Protein, hFc Tag Cat. No. PME100960



PRODUCT INFORMATION

SynonymsIFB;IFF;IFN-beta;IFNBDescriptionRecombinant human IFNB protein with C-terminal human FC tagDeliveryUnder developmentUniprot IDP01574Expression HostHEK293TagC-Human Fc TagMolecular CharacterizationIFNB(Met22-Asn187) hFc(Glu99-Ala330)Molecular WeightThe protein has a predicted molecular mass of 43.6 kDa after removal of the signal peptide. The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.Formulation & ReconstitutionLoophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. Storage & ShippingStorage & ShippingStorage 1.20°C to -80°C for 12 months in lyophilized from sterile protein signaling proteins, which are released as part of the innate immune response to pathogens. The protein screet of putty for ensering and thawing). Lyophilized proteins are shipped at ambient temperature. This gene encodes a cytokine that belongs to the interferons screetorin on is linked to autoimmune diseases. Mice deficient for this gene dispaling proteins, which are released as part of the innate immune response to pathogens. The protein encoded by this gene belongs to the type I interferons are involved in cell differentiation and anti-tumor defenses and chemokines. Overactivation of type I interferon secretion in response to a pathogen, type I interferons screetpor complex and induce transcription of genes such as those encoding inflammatory cytokines and chemokines. Overactivation of type I interferon. Interferon secretion on lisked to autoimmune diseases. Mice<	Target	IFNB
Descriptionhuman Fc tagDeliveryUnder developmentUniprot IDP01574Expression HostHEK293TagC-Human Fc TagMolecular CharacterizationIFNB(Met22-Asn187) hFc(Glu99-Ala330)Molecular WeightThe protein has a predicted molecular mass of 43.6 KD a dater removal of the signal peptide. The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining. Lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. Storage & ShippingStorage & ShippingStore at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized form ther interdence of the innate immune response to pathogens. The protein encoded by this gene belongs to the type I class of interferons, which are important for defense against viral infections. In addition, type I interferon serie involved in cell differentiation and anti-tumor defenses. Following secretion in response to a pathogen, type I interferons bind a homologous receptor complex and induce transcription of gene such as those encoding inflammatory cytokines and encoding inflammatory cytokines and 	Synonyms	IFB;IFF;IFN-beta;IFNB
Uniprof IDP01574Expression HostHEK293TagC-Human Fc TagMolecular CharacterizationIFNB(Met22-Asn187) hFc(Glu99-Ala330)Molecular WeightThe protein has a predicted molecular mass of 43.6 kDa after removal of the signal peptide. The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining. Lyophilization, Please see Certificate of Analysis for specific instructions of reconstitution. Storage & ShippingStorage & ShippingThis gene encodes a cytokine that belongs to the intended for use within a month, aliquot and store at -80° C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. This gene encodes a cytokine that belongs to the interferons sind of any defenses. Following secretion in response to a pathogen, type 1 interferons sare involved in cell differentiation and anti-tumor defenses. Following secretion in interferons are involved in cell differentiation and anti-tumor defenses. Following secretion in response to a pathogen, type 1 interferons sare involved in cell differentiation and anti-tumor defenses. Following secretion in interferons are involved in cell differentiation and anti-tumor defenses. Following secretion in interferons secretion. In addition. type 1 interferon secretion in treps ones to a phologen y cytokines and encoding inflammatory cytokines and encoding inflammatory cytokines and encluding defects in 8 cell maturation and increased susceptibility to viral infection. [provided by RefSeq, Sep 2015]UsageResearch use only	Description	
Expression HostHEK293TagC-Human Fc TagMolecular CharacterizationIFNB(Met22-Asn187) hFc(Glu99-Ala330)Molecular WeightThe protein has a predicted molecular mass of 43.6 kDa after removal of the signal peptide. The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining. Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilized from After reconstitution. Storage & ShippingStorage & ShippingExpression from the protein are shipped at ambient temperature. This gene encodes a cytokine that belongs to the interferon family of signaling proteins, which are released as part of the innate immune response to pathogens. The protein encoded by this gene belongs to the type 1 interferons are involved in cell differentiation and anti-tumor defenses. Following secretion in response to a pathogen, type 1 interferons and anti-tumor defenses. Following secretion in fergons end and the treferon secretion of type I interferon secretion of type I interferon secretion of type I interferon secretion of type I interferon secretion is finked to autoimmune diseases. Mice and induce transcription of genes such as those encoding inflammatory cytokines and chemokines. Overactivation of type I interferon secretion is inked to autoimmune diseases. Mice adefinited molecular secretion in the protection. [provided by RefSeq, Sep 2015]UsageResearch use only	Delivery	Under development
TagC-Human Fc TagMolecular CharacterizationIFNB(Met22-Asn187) hFc(Glu99-Ala330)Molecular WeightThe protein has a predicted molecular mass of 43.6 KDa after removal of the signal peptide. The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining. Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution. This gene encodes a cytokine that belongs to the interded for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. This gene encodes a cytokine that belongs to the interferon family of signaling proteins, which are released as part of the innate immune response to pathogens. The protein encoded by this gene belongs to the type I interferons are involved in cell differentiation and anti-tumor defenses against viral infections. In addition, type I interferon sector complex and induce transcription of genes such as those encoding inflammatory cytokines and chemokines. Overactivation of type I interferon secretion is linked to autoimmune diseases. Mice deficient for this gene display several phenotypes including defects in B cell maturation and increased susceptibility to viral infection. Iprovided by Refseq, Sep 2015]UsageResearch use only	Uniprot ID	P01574
Molecular CharacterizationIFNB(Met22-Asn187) hFc(Glu99-Ala330)Molecular WeightThe protein has a predicted molecular mass of 43.6 kDa after removal of the signal peptide. The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining. Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. Storage & ShippingStorage & Shipping	Expression Host	HEK293
CharacterizationIFNB(Met22-ASIL87) RFC(Glu99-Ala330)Molecular WeightThe protein has a predicted molecular mass of 43.6 kDa after removal of the signal peptide. The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining. Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. Storage & ShippingStorage & ShippingStorage at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. This gene encodes a cytokine that belongs to the interferon family of signaling proteins, which are released as part of the innate immume response to pathogens. The protein encoded by this gene belongs to the type I interferons are involved in cell differentiation and anti-tumor defenses. Following secretion in response to a pathogen, type I interferons for a homologous receptor complex and induce transcription of genes such as those encoding inflammatory cytokines and chemokines. Overactivation of type I interferon. Iprovided by RefSeq, Sep 2015]UsageResearch use only	Tag	C-Human Fc Tag
HoteCular weight43.6 kDa after removal of the signal peptide. The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining. Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilized form. After reconstitution. Storage & ShippingStorage & ShippingStorage at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.BackgroundBackgroundStorage as cytokine that belongs to the interferon family of signaling proteins, which are released as part of the innate immune response to pathogens. The protein encoded by this gene belongs to the type 1 class of interferons. In addition, type 1 interferons and induct proteins and to regulate the store and induct protein on of genes such as those encoding inflammatory cytokines and chemokines. Overactivation of type 1 interferon secretion in response to a pathogen, type 1 interferons and induce transcription of genes such as those encoding inflammatory cytokines and chemokines. Overactivation of type 1 interferon secretion in received by the year of the proteix and induce transcription of genes such as those encoding inflammatory cytokines and chemokines. Overactivation of type 1 interferon secretion in received by RefSeq, Sep 2015]BackgroundResearch use only		IFNB(Met22-Asn187) hFc(Glu99-Ala330)
Puritydetermined by SDS-PAGE and Coomassie blue staining. Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. Storage & ShippingStorage & ShippingStore at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.BackgroundThis gene encodes a cytokine that belongs to the interferon family of signaling proteins, which are released as part of the innate immune response to pathogens. The protein encoded by this gene belongs to the type I class of interferons, which are important for defense against viral infections. In addition, type I interferons are involved in cell differentiation and anti-tumor defenses. Following secretion in response to a pathogen, type I interferons bind a homologous receptor complex and induce transcription of genes such as those encoding inflammatory cytokines and chemokines. Overactivation of type I interferon secretion is linked to autoimmune diseases. Mice deficient for this gene display several phenotypes including defects in B cell maturation and increased susceptibility to viral infection. [provided by RefSeq, Sep 2015]UsageResearch use only	Molecular Weight	43.6 kDa after removal of the signal peptide.
Formulation & Reconstitution-8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.Background-8 month aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.Background-8 month aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.Background-8 month aliquot and store at induction, type I class of interferons, which are released as part of the innate immune response to pathogens. The protein encoded by this gene belongs to the type I class of interferons, which are important for defense against viral infections. In addition, type I interferons are involved in cell differentiation and anti-tumor defenses. Following secretion in response to a pathogen, type I interferons bind a homologous receptor complex and induce transcription of genes such as those encoding inflammatory cytokines and chemokines. Overactivation of type I interferon secretion is linked to autoimmune diseases. Mice deficient for this gene display several phenotypes including defects in B cell maturation and increased susceptibility to viral infection. [provided by RefSeq, Sep 2015]UsageResearch use only	Purity	determined by SDS-PAGE and Coomassie blue staining.
Storage & ShippingIyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. This gene encodes a cytokine that belongs to the interferon family of signaling proteins, which are released as part of the innate immune response to pathogens. The protein encoded by this gene belongs to the type I class of interferons, which are important for defense against viral infections. In addition, type I interferons are involved in cell differentiation and anti-tumor defenses. Following secretion in response to a pathogen, type I interferons bind a homologous receptor complex and induce transcription of genes such as those encoding inflammatory cytokines and chemokines. Overactivation of type I interferon secretion is linked to autoimmune diseases. Mice deficient for this gene display several phenotypes including defects in B cell maturation and increased susceptibility to viral infection. [provided by RefSeq, Sep 2015]UsageResearch use only		 - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
 Background Background interferon family of signaling proteins, which are released as part of the innate immune response to pathogens. The protein encoded by this gene belongs to the type I class of interferons, which are important for defense against viral infections. In addition, type I interferons are involved in cell differentiation and anti-tumor defenses. Following secretion in response to a pathogen, type I interferons bind a homologous receptor complex and induce transcription of genes such as those encoding inflammatory cytokines and chemokines. Overactivation of type I interferon secretion is linked to autoimmune diseases. Mice deficient for this gene display several phenotypes including defects in B cell maturation and increased susceptibility to viral infection. [provided by RefSeq, Sep 2015] Usage 	Storage & Shipping	lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient
	Background	interferon family of signaling proteins, which are released as part of the innate immune response to pathogens. The protein encoded by this gene belongs to the type I class of interferons, which are important for defense against viral infections. In addition, type I interferons are involved in cell differentiation and anti-tumor defenses. Following secretion in response to a pathogen, type I interferons bind a homologous receptor complex and induce transcription of genes such as those encoding inflammatory cytokines and chemokines. Overactivation of type I interferon secretion is linked to autoimmune diseases. Mice deficient for this gene display several phenotypes including defects in B cell maturation and increased susceptibility to viral infection.
Conjugate Unconjugated	Usage	
	Conjugate	Unconjugated

Email: info@dimabio.com Website: www.dimabio.com

