

PRODUCT INFORMATION

Target	IL-16
Synonyms	Pro-Interleukin-16; Interleukin-16; IL-16; Lymphocyte Chemoattractant Factor; LCF; IL16
Description	Recombinant Human Interleukin-16 is produced by our E.coli expression system and the target gene encoding Met1-Ser130 is expressed.
Delivery	In Stock
Uniprot ID	AAC12732.1
Expression Host	E.coli
Tag	
Molecular Characterization	Not available
Molecular Weight	13.4 KDa
Purity	Greater than 95% as determined by reducing SDS-PAGE.
Formulation & Reconstitution	Lyophilized from a 0.2 μ m filtered solution of 20mM PB, 150mM NaCl, pH 7.0. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Storage & Shipping	
Background	Interleukin-16 (IL-16) is a CD8 T cell-derived cytokine that induces chemotaxis of CD4 T cells and CD4 monocytes and eosinophils. Analysis by gel filtration suggests that, under physiological conditions, human IL-16 exists predominantly as a noncovalently linked multimer, but that some IL-16 may exist as a monomer. However, only the multimeric form appears to possess chemotactic activity, suggesting that receptor cross-linking may be required for activity. IL-16 also induces expression of IL-2 receptor (IL-2R) and MHC class II molecules on CD4 T cells. Human and murine IL-16 show significant cross-species reactivity.
Usage	Research use only
Conjugate	Unconjugated






Figure 1. Greater than 95% as determined by reducing SDS-PAGE.

DIMABIO CONFIDENTIAL

