

PRODUCT INFORMATION

| | |
|---|---|
| Target | IL1A |
| Synonyms | IL-1 alpha;IL-1A;IL1;IL1-ALPHA;IL1F1 |
| Description | Recombinant Human IL1A Protein with C-terminal human Fc tag |
| Delivery | In Stock |
| Uniprot ID | P01583 |
| Expression Host | HEK293 |
| Tag | C-Human Fc Tag |
| Molecular Characterization | IL1A(Ser113-Ala271) hFc(Glu99-Ala330) |
| Molecular Weight | The protein has a predicted molecular mass of 44.2 kDa after removal of the signal peptide. The apparent molecular mass of IL1A-hFc is approximately 35-55 kDa due to glycosylation. |
| Purity | The purity of the protein is greater than 90% as determined by SDS-PAGE and Coomassie blue staining. |
| Formulation & Reconstitution | Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. |
| Storage & Shipping | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Background | The protein encoded by this gene is a member of the interleukin 1 cytokine family. This cytokine is a pleiotropic cytokine involved in various immune responses, inflammatory processes, and hematopoiesis. This cytokine is produced by monocytes and macrophages as a proprotein, which is proteolytically processed and released in response to cell injury, and thus induces apoptosis. This gene and eight other interleukin 1 family genes form a cytokine gene cluster on chromosome 2. It has been suggested that the polymorphism of these genes is associated with rheumatoid arthritis and Alzheimer's disease. [provided by RefSeq, Jul 2008] |
| Usage | Research use only |






Figure 1. Human IL1A Protein, hFc Tag on SDS-PAGE under reducing condition.

Human IL1A, hFc Tagged protein ELISA

0.2 μ g of Human IL1A, hFc tagged protein per well




Figure 2. ELISA plate pre-coated by 2 μ g/mL (100 μ L/well) Human IL1A Protein, hFc Tag (PME101053) can bind Anti-IL1A Neutralizing antibody BME100097 in a linear range of 3.05-740.74 ng/mL.

