

PRODUCT INFORMATION

Target	IL36G
Synonyms	IL1E; IL1F9; IL1H1; IL-1F9; IL-1H1; IL1RP2; IL-1RP2
Description	Recombinant human IL36G Protein with C-terminal human Fc tag
Delivery	In Stock
Uniprot ID	Q9NZH8
Expression Host	HEK293
Tag	C-Human Fc tag
Molecular Characterization	IL36G(Ser18-Asp169) hFc(Glu99-Ala330)
Molecular Weight	The protein has a predicted molecular mass of 43.2 kDa after removal of the signal peptide.
Purity	The purity of the protein is greater than 90% as determined by SDS-PAGE and Coomassie blue staining.
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	The protein encoded by this gene is a member of the interleukin 1 cytokine family. The activity of this cytokine is mediated by interleukin 1 receptor-like 2 (IL1RL2/IL1R-rp2), and is specifically inhibited by interleukin 1 family, member 5 (IL1F5/IL-1 delta). Interferon-gamma, tumor necrosis factor-alpha and interleukin 1, beta (IL1B) are reported to stimulate the expression of this cytokine in keratinocytes. The expression of this cytokine in keratinocytes can also be induced by a contact hypersensitivity reaction or herpes simplex virus infection. This gene and eight other interleukin 1 family genes form a cytokine gene cluster on chromosome 2. [provided by RefSeq, May 2019]
Usage	Research use only
Conjugate	Unconjugated






Figure 1. Human IL36G Protein, hFc Tag on SDS-PAGE under reducing condition.

DIMABIO CONFIDENTIAL

