

PRODUCT INFORMATION

Tag C-Flag&Strep Tag

Target KCD11

Synonyms C17orf36, KCASH1, REN, REN/KCTD11

HEK293

Description Human KCD11-Strep full length protein-synthetic

Delivery 6~8weeks
Uniprot ID Q693B1

Protein Families Ion Channels: Other

Protein Pathways N/A

Expression Host

Background

Molecular Weight

The human full length KCD11-Strep protein has a

MW of 25.9 kDa

mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5%
– 8% trehalose is added as protectants before
lyophilization. Please see Certificate of Analysis
for specific instructions. Do not use solvents with

a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

Lyophilized from nanodisc solubilization buffer (20

Storage & Shipping intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

Plays a role as a marker and a regulator of neuronal differentiation; Up-regulated by a variety of neurogenic signals, such as retinoic acid, epidermal growth factor/EGF and

NGFB/nerve growth factor. Induces apoptosis, growth arrest and the expression of cyclindependent kinase inhibitor CDKN1B. Plays a role as a tumor repressor and inhibits cell growth and tumorigenicity of medulloblastoma (MDB). Acts as probable substrate-specific adapter for a BCR (BTB-CUL3-RBX1) E3 ubiquitin-protein ligase complex towards HDAC1. Functions as antagonist

complex towards HDAC1. Functions as antagonist of the Hedgehog pathway on cell proliferation and differentiation by affecting the nuclear transfer of transcription factor GLI1, thus maintaining cerebellar granule cells in undifferentiated state, this effect probably occurs via HDAC1 down-regulation, keeping GLI1 acetylated and inactive.

Email: info@dimabio.com Website: www.dimabio.com

When knock-down, Hedgehog antagonism is impaired and proliferation of granule cells is sustained. Activates the caspase

cascade.[UniProtKB/Swiss-Prot Function]

Usage Research use only
Conjugate Unconjugated



