

PRODUCT INFORMATION

|                              |   |
|------------------------------|---|
| Tag                          | C-Flag Tag  |
| Target                       | KCD17   |
| Synonyms                     | N/A   |
| Description                  | Human KCD17 full length protein-synthetic nanodisc  |
| Delivery                     | 6~8weeks  |
| Uniprot ID                   | Q8N5Z5  |
| Expression Host              | HEK293  |
| Protein Families             | Ion Channels: Other   |
| Protein Pathways             | N/A   |
| Molecular Weight             | The human full length KCD17 protein has a MW of 35.7kDa   |
| Formulation & Reconstitution | Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments.  |
| Storage & Shipping           | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.   |
| Background                   | This gene encodes a protein that belongs to a conserved family of potassium channel tetramerization domain (KCTD)-containing proteins. The encoded protein functions in ciliogenesis by acting as a substrate adaptor for the cullin3-based ubiquitin-conjugating enzyme E3 ligase, and targets trichoplein, a keratin-binding protein, for degradation via polyubiquitylation. A mutation in this gene is associated with autosomal dominant myoclonic dystonia 26. [provided by RefSeq, Nov 2016] |
| Usage                        | Research use only   |
| Conjugate                    | Unconjugated  |

