

PRODUCT INFORMATION

TargetKCTD7SynonymsCLN14, EPM3DescriptionHuman KCTD7-Strep full length protein-synthetic nanodiscDelivery6~8weeksUniprot IDQ96MP8Expression HostHEK293Protein FamiliesIon Channels: OtherProtein PathwaysN/AMolecular WeightThe human full length KCTD7-Strep protein has a MW of 33.1 kDa Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCI, 150 mM NaCI, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilized from noth solutions of divalent metal. Storage & ShippingStorage & ShippingStorage a ShippingAttract and the termerization domain-containing protein family. Family members are identified on a structural basis and contain a maino-terminal a structural basis and contain a maino-terminal
DescriptionHuman KCTD7-Strep full length protein-synthetic nanodiscDelivery6~8weeksUniprot IDQ96MP8Expression HostHEK293Protein FamiliesIon Channels: OtherProtein PathwaysN/AMolecular WeightThe human full length KCTD7-Strep protein has a MW of 33.1 kDa Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCI, 150 mM NaCI, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments.Storage & ShippingStorage & ShippingThis gene encodes a member of the potassium channel tetramerization domain-containing protein family. Family members are identified on a structural basis and contain an amino-terminal
DescriptionnanodiscnanodiscDelivery6~8weeksUniprot IDQ96MP8Expression HostHEK293Protein FamiliesIon Channels: OtherProtein PathwaysN/AMolecular WeightThe human full length KCTD7-Strep protein has a MW of 33.1 kDa Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCI, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. This gene encodes a member of the potassium channel tetramerization domain-containing protein family. Family members are identified on a structural basis and contain an amino-terminal
Uniprot IDQ96MP8Expression HostHEK293Protein FamiliesIon Channels: OtherProtein PathwaysN/AMolecular WeightThe human full length KCTD7-Strep protein has a MW of 33.1 kDa Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments. Storage & ShippingStorage & ShippingStorage e encodes a member of the potassium channel tetramerization domain-containing protein family. Family members are identified on a structural basis and contain an amino-terminal
Expression HostHEK293Protein FamiliesIon Channels: OtherProtein PathwaysN/AMolecular WeightThe human full length KCTD7-Strep protein has a MW of 33.1 kDa Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCI, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments. Storage & ShippingStorage & ShippingStorage encodes a member of the potassium channel tetramerization domain-containing protein family. Family members are identified on a structural basis and contain an amino-terminal
Protein FamiliesIon Channels: OtherProtein PathwaysN/AMolecular WeightThe human full length KCTD7-Strep protein has a MW of 33.1 kDa Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCI, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.Storage & ShippingThis gene encodes a member of the potassium channel tetramerization domain-containing protein family. Family members are identified on a structural basis and contain an amino-terminal
Protein PathwaysN/AMolecular WeightThe human full length KCTD7-Strep protein has a MW of 33.1 kDa Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments. Storage & ShippingStorage & ShippingStorage end -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized roteins are shipped at ambient temperature. This gene encodes a member of the potassium channel tetramerization domain-containing protein family. Family members are identified on a structural basis and contain an amino-terminal
Molecular WeightThe human full length KCTD7-Strep protein has a MW of 33.1 kDa Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. This gene encodes a member of the potassium channel tetramerization domain-containing protein family. Family members are identified on a structural basis and contain an amino-terminal
Molecular weightMW of 33.1 kDaMW of 33.1 kDaLyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments.Storage & ShippingStore at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.This gene encodes a member of the potassium channel tetramerization domain-containing protein family. Family members are identified on a structural basis and contain an amino-terminal
Formulation & ReconstitutionLyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments. Storage & ShippingStorage & ShippingStore at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. This gene encodes a member of the potassium channel tetramerization domain-containing protein family. Family members are identified on a structural basis and contain an amino-terminal
Storage & Shippingintended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.This gene encodes a member of the potassium channel tetramerization domain-containing protein family. Family members are identified on a structural basis and contain an amino-terminal
channel tetramerization domain-containing protein family. Family members are identified on a structural basis and contain an amino-terminal
Background domain similar to the T1 domain present in the voltage-gated potassium channel. Mutations in this gene have been associated with progressive myoclonic epilepsy-3. Alternative splicing results in multiple transcript variants.[provided by RefSeq, Jan 2011]
Usage Research use only
Conjugate Unconjugated

Email: info@dimabio.com Website: www.dimabio.com

