Description

Background



PRODUCT INFORMATION

KGF **Target**

Fibroblast growth factor 7;FGF-7;Heparin-binding growth factor 7;HBGF-7;Keratinocyte growth factor;FGF7 **Synonyms**

Recombinant Human Fibroblast Growth Factor 7/Keratinocyte growth factor is produced by our Mammalian expression system and the target gene encoding Cys32-Thr194 is expressed with a 6His tag at the C-terminus.

In Stock **Delivery Uniprot ID** P21781 **Expression Host HEK293** Tag C-6×His Tag

Molecular

Characterization **Molecular Weight** 20 KDa

Greater than 95% as determined by reducing **Purity**

SDS-PAGE.

Not available

Lyophilized from a 0.2 µm filtered solution of PBS, Formulation & Reconstitution

pH 7.4.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store Storage & Shipping

at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

témperature.

Fibroblast growth factor 7 (FGF7) is a secreted protein which is mainly located in epithelial cells and belongs to the heparin-binding growth factors family. FGF family members possess broad mitogenic and cell survival activities, and are involved in a variety of biological processes, including embryonic development, cell growth, morphogenesis, tissue repair, tumor growth and invasion. FGF7 is a potent epithelial cell-specific

growth factor, whose mitogenic activity is predominantly exhibited in keratinocytes but not in fibroblasts and endothelial cells. It is possible

major paracrine effector of normal epithelial cell

Email: info@dimabio.com Website: www.dimabio.com

proliferation.

Usage Research use only Conjugate Unconjugated








Figure 1. Greater than 95% as determined by reducing SDS-PAGE.




Figure 2. Measured in a cell proliferation assay using HaCaT cells. The ED50 for this effect is 10.94 ng/ml.



Email: info@dimabio.com Website: www.dimabio.com