

## **PRODUCT INFORMATION**

C-Flag&Strep Tag Tag

**Target KISSR** 

AXOR12; CPPB1; GPR54; HH8; HOT7T175; **Synonyms** 

KISS-1R

Human KISSR-Strep full length protein-synthetic Description

nanodisc

**Delivery** 6~8weeks **Uniprot ID** Q969F8 **HEK293 Expression Host** 

Formulation &

**Background** 

**Protein Families** Druggable Genome, GPCR, Transmembrane

**Protein Pathways** Neuroactive ligand-receptor interaction

The human full length KISSR-Strep protein has a **Molecular Weight** 

MW of 42.6 kDa

Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis

Reconstitution for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store

Storage & Shipping at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

The protein is a galanin-like G protein-coupled receptor that binds metastin, a peptide encoded by the metastasis suppressor gene KISS1. The

tissue distribution of the expressed gene suggests that it is involved in the regulation of

endocrine function, and this is supported by the finding that this gene appears to play a role in the onset of puberty. Mutations in this gene have been associated with hypogonadotropic

> Email: info@dimabio.com Website: www.dimabio.com

hypogonadism and central precocious puberty.

Usage Research use only

Conjugate Unconjugated

