

**PRODUCT INFORMATION**

|   |   |
|---|---|
| <b>Tag</b>                              | C-Flag&Strep Tag  |
| <b>Target</b>                           | KISSR   |
| <b>Synonyms</b>                         | AXOR12; CPPB1; GPR54; HH8; HOT7T175; KISS-1R  |
| <b>Description</b>                      | Human KISSR-Strep full length protein-synthetic nanodisc  |
| <b>Delivery</b>                         | 6~8weeks  |
| <b>Uniprot ID</b>                       | Q969F8  |
| <b>Expression Host</b>                  | HEK293  |
| <b>Protein Families</b>                 | Druggable Genome, GPCR, Transmembrane   |
| <b>Protein Pathways</b>                 | Neuroactive ligand-receptor interaction   |
| <b>Molecular Weight</b>                 | The human full length KISSR-Strep protein has a MW of 42.6 kDa  |
| <b>Formulation &amp; Reconstitution</b> | Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for   |
| <b>Storage &amp; Shipping</b>           | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.   |
| <b>Background</b>                       | The protein is a galanin-like G protein-coupled receptor that binds metastin, a peptide encoded by the metastasis suppressor gene KISS1. The tissue distribution of the expressed gene suggests that it is involved in the regulation of endocrine function, and this is supported by the finding that this gene appears to play a role in the onset of puberty. Mutations in this gene have been associated with hypogonadotropic hypogonadism and central precocious puberty. |
| <b>Usage</b>                            | Research use only   |
| <b>Conjugate</b>                        | Unconjugated  |

