

PRODUCT INFORMATION

Target	M-CSF
Synonyms	Macrophage Colony-Stimulating Factor 1;CSF-1;M-CSF;MCSF;Lanimoestim;CSF1
Description	Recombinant Human Macrophage Colony-Stimulating Factor is produced by our Mammalian expression system and the target gene encoding Glu33-Arg255 is expressed with a 6His tag at the C-terminus.
Delivery	In Stock
Uniprot ID	P09603
Expression Host	HEK293
Tag	C-6×His Tag
Molecular Characterization	Not available
Molecular Weight	26.17 KDa
Purity	Greater than 95% as determined by reducing SDS-PAGE.
Formulation & Reconstitution	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	Macrophage Colony-Stimulating Factors (m-csf) are cytokines that act in hematopoiesis by controlling the production, differentiation, and function of 2 related white cell populations of the blood, the granulocytes and the monocytes-macrophages. CSF-1 promotes the release of proinflammatory chemokines, and thereby plays an important role in innate immunity and in inflammatory processes. It also plays an important role in the regulation of osteoclast proliferation and differentiation, the regulation of bone resorption, and is required for normal bone development. CSF-1 is required for normal male and female fertility and promotes reorganization of the actin cytoskeleton, regulates formation of membrane ruffles, cell adhesion and cell migration. It also plays a role in lipoprotein clearance.
Usage	Research use only
Conjugate	Unconjugated






Figure 1. Greater than 95% as determined by reducing SDS-PAGE.

