

PRODUCT INFORMATION

Tag	N-MBP Tag, C-Flag Tag
Target	AGTR1
Synonyms	AG2S; AGTR1B; AT1; AT1AR; AT1B; AT1BR; AT1R; AT2R1; HAT1R
Description	Human MBP-AGTR1 full length protein-synthetic nanodisc
Delivery	In Stock
Uniprot ID	P30556
Expression Host	HEK293
Protein Families	Druggable Genome, GPCR, Transmembrane
Protein Pathways	Calcium signaling pathway, Neuroactive ligand-receptor interaction, Renin-angiotensin system, Vascular smooth muscle contraction
Molecular Weight	The human full length MBP-AGTR1 protein has a MW of 81.1 kDa Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments.
Formulation & Reconstitution	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Storage & Shipping	Angiotensin II is a potent vasopressor hormone and a primary regulator of aldosterone secretion. It is an important effector controlling blood pressure and volume in the cardiovascular system. It acts through at least two types of receptors. This gene encodes the type 1 receptor which is thought to mediate the major cardiovascular effects of angiotensin II. This gene may play a role in the generation of reperfusion arrhythmias following restoration of blood flow to ischemic or infarcted myocardium. It was previously thought that a related gene, denoted as AGTR1B, existed; however, it is now believed that there is only one type 1 receptor gene in humans. Alternative splicing of this gene results in multiple transcript variants.
Background	
Usage	Research use only
Conjugate	Unconjugated



ELISA assay to evaluate MBP-AGTR1-Nanodisc
0.2µg Human MBP-AGTR1-Nanodisc per well




Figure 1. Elisa plates were pre-coated with N-MBP Tag, C-Flag Tag MBP-AGTR1-Nanodisc (0.2µg/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with MBP-AGTR1-Nanodisc is 4.309ng/ml.



Figure 2. Human MBP-AGTR1-Nanodisc with N-MBP Tag, C-Flag Tag on SDS-PAGE

