

## **PRODUCT INFORMATION**

Тад	C-Flag&Strep Tag
Target	MFSD13A
Synonyms	bA18I14.8; C10orf77; TMEM180
Description	Human MFSD13A-Strep full length protein- synthetic nanodisc
Delivery	6~8weeks
Uniprot ID	Q14CX5
<b>Expression Host</b>	HEK293
<b>Protein Families</b>	Transmembrane
Protein Pathways	N/A
Molecular Weight	The human full length MFSD13A-Strep protein has a MW of 57.4 kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	MFSD13A, also called Transmembrane protein 180 (TMEM180), is a transmembrane protein that belongs to the glycoside-pentoside-hexuronide (GPH):cation symporter family. Members of this family catalyze symport of a sugar molecule with a monovalent cation (H or Na ). MFSD13A is classified as a member of the cation symporter family and a multi-pass membrane protein, but little information is available regarding its substrate and topology.
Usage	Research use only
Conjugate	Unconjugated

