

PRODUCT INFORMATION

| | |
|---|--|
| Target | MOG |
| Synonyms | BTN6; BTNL11; MOGIG2; NRCLP7 |
| Description | Recombinant human MOG Protein with C-terminal human Fc tag |
| Delivery | In Stock |
| Uniprot ID | Q16653 |
| Expression Host | HEK293 |
| Tag | C-Human Fc tag |
| Molecular Characterization | MOG(Gly30-Gly154) hFc(Glu99-Ala330) |
| Molecular Weight | The protein has a predicted molecular mass of 40.4 kDa after removal of the signal peptide. |
| Purity | The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining. |
| Formulation & Reconstitution | Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. |
| Storage & Shipping | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Background | The product of this gene is a membrane protein expressed on the oligodendrocyte cell surface and the outermost surface of myelin sheaths. Due to this localization, it is a primary target antigen involved in immune-mediated demyelination. This protein may be involved in completion and maintenance of the myelin sheath and in cell-cell communication. Alternatively spliced transcript variants encoding different isoforms have been identified. [provided by RefSeq, Jul 2008] |
| Usage | Research use only |
| Conjugate | Unconjugated |






Figure 1. Human MOG Protein, hFc Tag on SDS-PAGE under reducing condition.

DIMABIO CONFIDENTIAL

