

PRODUCT INFORMATION

| | |
|---|---|
| Tag | C-Flag&Strep Tag |
| Target | MTR1A |
| Synonyms | MEL-1A-R, MT1 |
| Description | Human MTR1A-Strep full length protein-synthetic nanodisc |
| Delivery | 6~8weeks |
| Uniprot ID | P48039 |
| Expression Host | HEK293 |
| Protein Families | GPCR,Transmembrane,Druggable Genome, |
| Protein Pathways | GPCRDB Class A Rhodopsin-like,Small ligand GPCRs,Cancer, |
| Molecular Weight | The human full length MTR1A-Strep protein has a MW of 39.4 kDa Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments. |
| Formulation & Reconstitution | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Storage & Shipping | |
| Background | This gene encodes one of two high affinity forms of a receptor for melatonin, the primary hormone secreted by the pineal gland. This receptor is a G-protein coupled, 7-transmembrane receptor that is responsible for melatonin effects on mammalian circadian rhythm and reproductive alterations affected by day length. The receptor is an integral membrane protein that is readily detectable and localized to two specific regions of the brain. The hypothalamic suprachiasmatic nucleus appears to be involved in circadian rhythm while the hypophysial pars tuberalis may be responsible for the reproductive effects of melatonin. [provided by RefSeq, Jul 2008] |
| Usage | Research use only |
| Conjugate | Unconjugated |

