

PRODUCT INFORMATION

C-Flag&Strep Tag Tag

Target NK2R

Synonyms NKNAR; SKR; TAC2R

Human NK2R-Strep full length protein-synthetic **Description**

nanodisc

Delivery In Stock **Uniprot ID** P21452 **Expression Host HEK293**

Formulation &

Reconstitution

Background

Protein Families Druggable Genome, GPCR, Transmembrane

Calcium signaling pathway, Neuroactive ligand-**Protein Pathways**

receptor interaction

The human full length NK2R-Strep protein has a **Molecular Weight**

MW of 44.4 kDa

Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis

for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater

than 5 mM) in subsequent experiments. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store Storage & Shipping at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

This protein belongs to a family of proteins that function as receptors for tachykinins. Receptor affinities are specified by variations in the 5'-end

of the sequence. The receptors belonging to this family are characterized by interactions with G proteins and 7 hydrophobic transmembrane

regions.

Usage Research use only

Conjugate Unconjugated



Email: info@dimabio.com Website: www.dimabio.com





ELISA assay to evaluate NK2R-Strep-Nanodisc 0.2µg Human NK2R-Strep-Nanodisc per well




Figure 1. Elisa plates were pre-coated with C-Flag&Strep Tag NK2R-Strep-Nanodisc (0.2 μ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with NK2R-Strep-nanodisc is 2.571ng/ml.




Figure 2. Human NK2R-Strep-Nanodisc, C-Flag&Strep Tag on SDS-PAGE

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

Email: info@dimabio.com Website: www.dimabio.com

