

PRODUCT INFORMATION

Tag	C-Flag&Strep Tag
Target	NK3R
Synonyms	HH11, NK-3R, NK3, NK3R, NKR, TAC3R, TAC3RL
Description	Human NK3R-Strep full length protein-synthetic nanodisc
Delivery	6~8weeks
Uniprot ID	P29371
Expression Host	HEK293
Protein Families	Transmembrane,Druggable Genome,
Protein Pathways	Peptide GPCRs,
Molecular Weight	The human full length NK3R-Strep protein has a MW of 52.2 kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	This gene belongs to a family of genes that function as receptors for tachykinins. Receptor affinities are specified by variations in the 5'-end of the sequence. The receptors belonging to this family are characterized by interactions with G proteins and 7 hydrophobic transmembrane regions. This gene encodes the receptor for the tachykinin neurokinin 3, also referred to as neurokinin B. [provided by RefSeq, Jul 2008]
Usage	Research use only
Conjugate	Unconjugated

