

PRODUCT INFORMATION

| | |
|---|--|
| Tag | C-Flag Tag |
| Target | NMD3A |
| Synonyms | GluN3A, NMDAR-L, NMDAR3A, NR3A |
| Description | Human NMD3A full length protein-synthetic nanodisc |
| Delivery | 6~8weeks |
| Uniprot ID | Q8TCU5 |
| Expression Host | HEK293 |
| Protein Families | Ion Channels: Glutamate Receptors |
| Protein Pathways | N/A |
| Molecular Weight | The human full length NMD3A protein has a MW of 125.5kDa |
| Formulation & Reconstitution | Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments. |
| Storage & Shipping | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Background | This gene encodes a subunit of the N-methyl-D-aspartate (NMDA) receptors, which belong to the superfamily of glutamate-regulated ion channels, and function in physiological and pathological processes in the central nervous system. This subunit shows greater than 90% identity to the corresponding subunit in rat. Studies in the knockout mouse deficient in this subunit suggest that this gene may be involved in the development of synaptic elements by modulating NMDA receptor activity. [provided by RefSeq, Jul 2008] |
| Usage | Research use only |
| Conjugate | Unconjugated |

