

PRODUCT INFORMATION

Tag	C-Flag&Strep Tag
Target	NMD3A
Synonyms	GluN3A, NMDAR-L, NMDAR3A, NR3A
Description	Human NMD3A-Strep full length protein-synthetic nanodisc
Delivery	6~8weeks
Uniprot ID	Q8TCU5
Expression Host	HEK293
Protein Families	Ion Channels: Glutamate Receptors
Protein Pathways	N/A
Molecular Weight	The human full length NMD3A-Strep protein has a MW of 125.5 kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	This gene encodes a subunit of the N-methyl-D-aspartate (NMDA) receptors, which belong to the superfamily of glutamate-regulated ion channels, and function in physiological and pathological processes in the central nervous system. This subunit shows greater than 90% identity to the corresponding subunit in rat. Studies in the knockout mouse deficient in this subunit suggest that this gene may be involved in the development of synaptic elements by modulating NMDA receptor activity. [provided by RefSeq, Jul 2008]
Usage	Research use only
Conjugate	Unconjugated

