

PRODUCT INFORMATION

Tag	C-Flag Tag
Target	NPY1R
Synonyms	NPY1-R; NPYR
Description	Human NPY1R full length protein-synthetic nanodisc
Delivery	In Stock
Uniprot ID	P25929
Expression Host	HEK293
Protein Families	Druggable Genome, GPCR, Transmembrane
Protein Pathways	Neuroactive ligand-receptor interaction
Molecular Weight	The human full length NPY1R protein has a MW of 44.4 kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	The transmembrane protein mediates the function of neuropeptide Y (NPY), a neurotransmitter, and peptide YY (PYY), a gastrointestinal hormone. The encoded receptor undergoes fast agonist-induced internalization through clathrin-coated pits and is subsequently recycled back to the cell membrane. Activation of Y1 receptors may result in mobilization of intracellular calcium and inhibition of adenylate cyclase activity.
Usage	Research use only
Conjugate	Unconjugated



ELISA assay to evaluate NPY1R-Nanodisc 0.2 μ g Human NPY1R-Nanodisc per well




Figure 1. Elisa plates were pre-coated with Flag Tag NPY1R-Nanodisc (0.2 μ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with NPY1R-Nanodisc is 5.641ng/ml.

kDa M R

250
130
100
70
55
35
25
15
10



Figure 2. WB analysis of Human NPY1R-Nanodisc with anti-Flag monoclonal antibody at 1/5000 dilution, followed by Goat Anti-Rabbit IgG HRP at 1/5000 dilution

