

**PRODUCT INFORMATION**

|   |  |
|---|--|
| <b>Tag</b>                              | C-Flag&Strep Tag   |
| <b>Target</b>                           | OR2F1  |
| <b>Synonyms</b>                         | 7M1-2, OLF3, OR14-60, OR2F3, OR2F3P, OR2F4, OR2F5, OR7-139, OR7-140  |
| <b>Description</b>                      | Human OR2F1-Strep full length protein-synthetic nanodisc   |
| <b>Delivery</b>                         | 6~8weeks   |
| <b>Uniprot ID</b>                       | Q13607   |
| <b>Expression Host</b>                  | HEK293   |
| <b>Protein Families</b>                 | Transmembrane,Druggable Genome,  |
| <b>Protein Pathways</b>                 | GPCRDB Class A Rhodopsin-like,GPCRDB Other,  |
| <b>Molecular Weight</b>                 | The human full length OR2F1-Strep protein has a MW of 35.4 kDa<br>Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments.   |
| <b>Formulation &amp; Reconstitution</b> | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.  |
| <b>Storage &amp; Shipping</b>           |  |
| <b>Background</b>                       | Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. This olfactory receptor gene is a segregating pseudogene, where some individuals have an allele that encodes a functional olfactory receptor, while other individuals have an allele encoding a protein that is predicted to be non-functional. [provided by RefSeq, Jun 2015] |
| <b>Usage</b>                            | Research use only  |
| <b>Conjugate</b>                        | Unconjugated   |

