Cat. No. FLP100429

Delivery

Background



PRODUCT INFORMATION

C-Flag Tag Tag

OXYR Target

OT-R; OTR **Synonyms**

Human OXYR full length protein-synthetic **Description**

nanodisc In Stock

Uniprot ID P30559 **Expression Host HEK293**

Protein Families Druggable Genome, GPCR, Transmembrane

Calcium signaling pathway, Neuroactive ligand-**Protein Pathways**

receptor interaction

The human full length OXYR protein has a MW of **Molecular Weight**

42.8 kDa

Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis

Formulation & Reconstitution for specific instructions. Do not use solvents with

a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store Storage & Shipping at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

The protein belongs to the G-protein coupled receptor family and acts as a receptor for oxytocin. Its activity is mediated by G proteins

which activate a phosphatidylinositol-calcium second messenger system. The oxytocin-oxytocin receptor system plays an important role in the

Email: info@dimabio.com Website: www.dimabio.com

uterus during parturition.

Usage Research use only Conjugate Unconjugated





ELISA assay to evaluate OXYR-Nanodisc 0.2μg Human OXYR-Nanodisc per well




Figure 1. Elisa plates were pre-coated with Flag Tag OXYR-Nanodisc (0.2 μ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with OXYR-Nanodisc is 11.52ng/ml.




Figure 2. Human OXYR-Nanodisc, Flag Tag on SDS-PAGE

Email: info@dimabio.com
Website: www.dimabio.com

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)