

PRODUCT INFORMATION

Tag	C-Flag&Strep Tag
Target	P2RX7
Synonyms	P2X7
Description	Human P2RX7-Strep full length protein-synthetic nanodisc
Delivery	6~8weeks
Uniprot ID	Q99572
Expression Host	HEK293
Protein Families	Druggable Genome, Ion Channels: ATP Receptors, Transmembrane
Protein Pathways	Calcium signaling pathway, Neuroactive ligand-receptor interaction
Molecular Weight	The human full length P2RX7-Strep protein has a MW of 68.4 kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	The product of this gene belongs to the family of purinoceptors for ATP. This receptor functions as a ligand-gated ion channel and is responsible for ATP-dependent lysis of macrophages through the formation of membrane pores permeable to large molecules. Activation of this nuclear receptor by ATP in the cytoplasm may be a mechanism by which cellular activity can be coupled to changes in gene expression. Multiple alternatively spliced variants have been identified, most of which fit nonsense-mediated decay (NMD) criteria. [provided by RefSeq, Jul 2010]
Usage	Research use only
Conjugate	Unconjugated

