

PRODUCT INFORMATION

PDL1 **Target**

PD-L1;CD274;B7-H1;PDCD1L1;PDCD1LG1 **Synonyms** Recombinant human PD-L1 protein with C-**Description**

terminal 6×His tag

Delivery In Stock **Uniprot ID** Q9NZQ7 **Expression Host HEK293** Tag C-6×His Tag

Molecular

Background

PD-L1(Phe19-Arg238) 6×His tag Characterization

The protein has a predicted molecular mass of

26.0 kDa after removal of the signal peptide. The apparent molecular mass of PDL1-His is **Molecular Weight**

approximately 33-35 kDa due to glycosylation.

The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue Purity

staining.

Lyophilized from sterile PBS, pH 7.4. Normally 5 % 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Formulation & Reconstitution

for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store

Storage & Shipping at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

This gene encodes an immune inhibitory receptor ligand that is expressed by hematopoietic and non-hematopoietic cells, such as T cells and B cells and various types of tumor cells. The encoded protein is a type I transmembrane protein that has immunoglobulin V-like and C-like domains. Interaction of this ligand with its receptor inhibits T-cell activation and cytokine production. During infection or inflammation of normal tissue, this interaction is important for

preventing autoimmunity by maintaining homeostasis of the immune response. In tumor microenvironments, this interaction provides an immune escape for tumor cells through cytotoxic T-cell inactivation. Expression of this gene in tumor cells is considered to be prognostic in many types of human malignancies, including colon cancer and renal cell carcinoma. Alternative

> Email: info@dimabio.com Website: www.dimabio.com

splicing results in multiple transcript variants.

Usage Research use only Conjugate Unconjugated

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China)






Figure 1. Human PD-L1 Protein, His Tag on SDS-PAGE under reducing condition.



