

PRODUCT INFORMATION

| Тад | C-Flag Tag |
|---------------------------------|---|
| Target | PK2L1 |
| Synonyms | PCL, PKD2L, PKDL, TRPP3 |
| Description | Human PK2L1 full length protein-synthetic nanodisc |
| Delivery | 6~8weeks |
| Uniprot ID | Q9P0L9 |
| Expression Host | HEK293 |
| Protein Families | Ion Channels: Transient receptor potential |
| Protein Pathways | N/A |
| Molecular Weight | The human full length PK2L1 protein has a MW of 92kDa |
| Formulation & Reconstitution | Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments. Store at -20°C to -80°C for 12 months in |
| Storage & Shipping | lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Background | This gene encodes a member of the polycystin protein family. The encoded protein contains multiple transmembrane domains, and cytoplasmic N- and C-termini. The protein may be an integral membrane protein involved in cell- cell/matrix interactions. This protein functions as a calcium-regulated nonselective cation channel. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Dec 2011] |
| Usage | Research use only |
| Conjugate | Unconjugated |
| | |

