

PRODUCT INFORMATION

| | |
|---|---|
| Target | POSTN |
| Synonyms | OSF-2;OSF2;PDLPOSTN;PN |
| Description | Recombinant Human POSTN Protein with C-terminal 6×His tag |
| Delivery | In Stock |
| Uniprot ID | Q15063 |
| Expression Host | HEK293 |
| Tag | C-6×His Tag |
| Molecular Characterization | POSTN(Asn22-Gln836) 6×His tag |
| Molecular Weight | The protein has a predicted molecular mass of 91.8 kDa after removal of the signal peptide. The apparent molecular mass of POSTN-His is approximately 55-100 kDa due to glycosylation. |
| Purity | The purity of the protein is greater than 85% as determined by SDS-PAGE and Coomassie blue staining. |
| Formulation & Reconstitution | Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. |
| Storage & Shipping | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Background | This gene encodes a secreted extracellular matrix protein that functions in tissue development and regeneration, including wound healing, and ventricular remodeling following myocardial infarction. The encoded protein binds to integrins to support adhesion and migration of epithelial cells. This protein plays a role in cancer stem cell maintenance and metastasis. Mice lacking this gene exhibit cardiac valve disease, and skeletal and dental defects. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Sep 2015] |
| Usage | Research use only |






Figure 1. Human POSTN Protein, His Tag on SDS-PAGE under reducing condition.

DIMABIO CONFIDENTIAL

