

## **PRODUCT INFORMATION**

C-Flag&Strep Tag Tag

**Target** S1PR1

CD363, CHEDG1, D1S3362, ECGF1, EDG-1, EDG1, **Synonyms** 

S1P1

Human S1PR1-Strep full length protein-synthetic Description

nanodisc

**Delivery** 6~8weeks **Uniprot ID** P21453 **Expression Host HEK293** 

Formulation &

Reconstitution

**Background** 

Storage & Shipping

**Protein Families** GPCR, Transmembrane, Druggable Genome,

GPCRDB Other, S1P Signaling, Small ligand GPCRs,Angiogenesis,Angiogenesis,G-Protein Coupled Receptors Signaling Pathway, **Protein Pathways** 

The human full length S1PR1-Strep protein has a **Molecular Weight** 

MW of 42.8 kDa

Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with

a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments. Store at -20°C to -80°C for 12 months in

lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient

témperature.

The protein encoded by this gene is structurally similar to G protein-coupled receptors and is highly expressed in endothelial cells. It binds the ligand sphingosine-1-phosphate with high affinity and high specificity, and suggested to be involved in the processes that regulate the differentiation

of endothelial cells. Activation of this receptor induces cell-cell adhesion. Alternative splicing results in multiple transcript variants. [provided

> Email: info@dimabio.com Website: www.dimabio.com

by RefSeq, Mar 2016]

**Usage** Research use only

Conjugate Unconjugated



/+86-400-006-0995(China)

