

**PRODUCT INFORMATION**

<b>Target</b>	SLC25A4
<b>Synonyms</b>	AAC1; ANT; ANT 1; ANT1; MTDPS12; MTDPS12A; PEO2; PEO3; PEOA2; T1
<b>Description</b>	Human SLC25A4 full length protein-MNP
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	P12235
<b>Expression Host</b>	HEK293
<b>Protein Families</b>	Druggable Genome, Transmembrane
<b>Protein Pathways</b>	Calcium signaling pathway, Huntington's disease, Parkinson's disease
<b>Molecular Weight</b>	The human full length SLC25A4 protein has a MW of 33.1 kDa
<b>Formulation &amp; Reconstitution</b>	Lyophilized from PBS. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions.
<b>Storage &amp; Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	The protein functions as a gated pore that translocates ADP from the cytoplasm into the mitochondrial matrix and ATP from the mitochondrial matrix into the cytoplasm. The protein forms a homodimer embedded in the inner mitochondria membrane. Mutations in this gene have been shown to result in autosomal dominant progressive external ophthalmoplegia and familial hypertrophic cardiomyopathy.
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated



### ELISA assay to evaluate SLC25A4-MNP 0.5 $\mu$ g Human SLC25A4-MNP per well




Figure1. Elisa plates were pre-coated with 0.5 $\mu$ g/per well purified human SLC25A4 full length membrane nanoparticles. Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with SLC25A4 full length membrane nanoparticles is 44.02ng/ml.

