

PRODUCT INFORMATION

Тад	C-Flag Tag
Target	SSR1
Synonyms	SRIF-2, SS-1-R, SS1-R, SS1R
Description	Human SSR1 full length protein-synthetic nanodisc
Delivery	6~8weeks
Uniprot ID	P30872
Expression Host	HEK293
Protein Families	GPCR,Transmembrane,Druggable Genome,
Protein Pathways	GPCRDB Class A Rhodopsin-like,Peptide GPCRs,Cancer,
Molecular Weight	The human full length SSR1 protein has a MW of 42.7kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments. Store at -20°C to -80°C for 12 months in
Storage & Shipping	lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	Somatostatins are peptide hormones that regulate diverse cellular functions such as neurotransmission, cell proliferation, and endocrine signaling as well as inhibiting the release of many hormones and other secretory proteins. Somatostatin has two active forms of 14 and 28 amino acids. The biological effects of somatostatins are mediated by a family of G- protein coupled somatostatin receptors that are expressed in a tissue-specific manner. The protein encoded by this gene is a member of the superfamily of somatostatin receptors having seven transmembrane segments. Somatostatin receptors form homodimers and heterodimers with other members of the superfamily as well as with other G-protein coupled receptors and receptor tyrosine kinases. This somatostatin-14 than -28. [provided by RefSeq, Jul 2012]
Usage	Research use only
Conjugate	Unconjugated

Email: info@dimabio.com Website: www.dimabio.com

