

PRODUCT INFORMATION

| | |
|---|--|
| Tag | C-Flag Tag |
| Target | TRPC5 |
| Synonyms | PPP1R159, TRP5 |
| Description | Human TRPC5 full length protein-synthetic nanodisc |
| Delivery | 6~8weeks |
| Uniprot ID | Q9UL62 |
| Expression Host | HEK293 |
| Protein Families | Ion Channels: Transient receptor potential |
| Protein Pathways | N/A |
| Molecular Weight | The human full length TRPC5 protein has a MW of 111.4kDa |
| Formulation & Reconstitution | Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments. |
| Storage & Shipping | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Background | This gene belongs to the transient receptor family. It encodes one of the seven mammalian TRPC (transient receptor potential channel) proteins. The encoded protein is a multi-pass membrane protein and is thought to form a receptor-activated non-selective calcium permeant cation channel. The protein is active alone or as a heteromultimeric assembly with TRPC1, TRPC3, and TRPC4. It also interacts with multiple proteins including calmodulin, CABP1, enkurin, Na(+)-H+ exchange regulatory factor (NHERF), interferon-induced GTP-binding protein (MX1), ring finger protein 24 (RNF24), and SEC14 domain and spectrin repeat-containing protein 1 (SESTD1). [provided by RefSeq, May 2010] |
| Usage | Research use only |
| Conjugate | Unconjugated |

