

PRODUCT INFORMATION

Tag C-Flag Tag TRPC5 **Target**

Synonyms PPP1R159, TRP5

Human TRPC5 full length protein-synthetic **Description**

nanodisc **Delivery** 6~8weeks **Uniprot ID** Q9UL62 **Expression Host HEK293**

Protein Families Ion Channels: Transient receptor potential

Protein Pathways N/A

Storage & Shipping

Background

Usage

The human full length TRPC5 protein has a MW of **Molecular Weight**

111.4kDa

mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Formulation & Reconstitution for specific instructions. Do not use solvents with

a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

Lyophilized from nanodisc solubilization buffer (20

intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient

témperature.

This gene belongs to the transient receptor family, It encodes one of the seven mammalian TRPC (transient receptor potential channel) proteins. The encoded protein is a multi-pass membrane protein and is thought to form a receptor-activated non-selective calcium permeant cation channel. The protein is active alone or as a heteromultimeric assembly with TRPC1, TRPC3, and TRPC4. It also interacts with

multiple proteins including calmodulin, CABP1, enkurin, Na(+)-H+ exchange regulatory factor (NHERF), interferon-induced GTP-binding protein (MX1), ring finger protein 24 (RP24), and SEC14 domain and spectrin repeat-containing protein 1 (SESTD1). [provided by RefSeq, May 2010]

> Email: info@dimabio.com Website: www.dimabio.com

Research use only

Conjugate Unconjugated

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China)

/+86-400-006-0995(China)

