

PRODUCT INFORMATION

Tag	C-Flag&Strep Tag
Target	TRPM7
Synonyms	ALSPDC, CHAK, CHAK1, LTRPC7, LTrpC-7, TRP-PLIK
Description	Human TRPM7-Strep full length protein-synthetic nanodisc
Delivery	6~8weeks
Uniprot ID	Q96QT4
Expression Host	HEK293
Protein Families	Ion Channels: Transient receptor potential
Protein Pathways	N/A
Molecular Weight	The human full length TRPM7-Strep protein has a MW of 212.7 kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	This gene belongs to the melastatin subfamily of transient receptor potential family of ion channels. The protein encoded by this gene is both an ion channel and a serine/threonine protein kinase. The kinase activity is essential for the ion channel function, which serves to increase intracellular calcium levels and to help regulate magnesium ion homeostasis. The encoded protein is involved in cytoskeletal organization, cell adhesion, cell migration and organogenesis. Defects in this gene are a cause of amyotrophic lateral sclerosis-parkinsonism/dementia complex of Guam. The gene may also be associated with defects of cardiac function. [provided by RefSeq, Aug 2017]
Usage	Research use only
Conjugate	Unconjugated

