

PRODUCT INFORMATION

Target	TSPAN33
Synonyms	PEN; PEN.; TSPAN-33
Description	Human TSPAN33 full length protein-synthetic nanodisc
Delivery	In Stock
Uniprot ID	Q86UF1
Expression Host	HEK293
Protein Families	Transmembrane
Protein Pathways	N/A
Molecular Weight	The human full length TSPAN33 protein has a MW of 31.5 kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with pH lower than 6.5 in subsequent experiments.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	Plays an important role in normal erythropoiesis (By similarity). It has a role in the differentiation of erythroid progenitors (By similarity). Regulates maturation and trafficking of the transmembrane metalloprotease ADAM10 (PubMed:26686862). Negatively regulates ligand-induced Notch activity probably by regulating ADAM10 activity (PubMed:26686862).
Usage	Research use only



ELISA assay to evaluate TSPAN33-Nanodisc 0.2 μ g Human TSPAN33-Nanodisc per well




Figure1. Elisa plates were pre-coated with Flag Tag TSPAN33-Nanodisc (0.2 μ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with TSPAN33-Nanodisc is 10.31ng/ml.




Figure2. Human TSPAN33-Nanodisc, Flag Tag on SDS-PAGE

