

PRODUCT INFORMATION

C-Flag&Strep Tag Tag

VIPR2 **Target**

C16DUPq36.3, DUP7q36.3, PACAP-R-3, PACAP-R3, VIP-R-2, VPAC2, VPAC2R, VPCAP2R **Synonyms**

Human VIPR2-Strep full length protein-synthetic Description

nanodisc

Delivery 6~8weeks **Uniprot ID** P41587 **HEK293 Expression Host**

Formulation &

Reconstitution

Background

Protein Families GPCR, Transmembrane, Druggable Genome,

GPCRDB Class B Secretin-like, **Protein Pathways**

The human full length VIPR2-Strep protein has a **Molecular Weight**

MW of 49.5 kDa

Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with

a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store

Storage & Shipping at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

This gene encodes a receptor for vasoactive intestinal peptide, a small neuropeptide. Vasoactive intestinal peptide is involved in smooth muscle relaxation, exocrine and endocrine secretion, and water and ion flux in lung and intestinal epithelia. Its actions are

effected through integral membrane receptors associated with a guanine nucleotide binding protein which activates adenylate cyclase.

[provided by RefSeq, Aug 2011]

Usage Research use only Conjugate Unconjugated



